arXiv:1205.6460v1 [math.CO] 29 May 2012

A Combinatorial Approach to Positional Number Systems

Andrew Vince
Department of Mathematics, University of Florida
Gainesville, FL, USA
`avince@ufl.edu`

Abstract

Although the representation of the real numbers in terms of a base and a set of digits has a long history, new questions arise even in simple situations. This paper concerns binary radix systems, i.e., positional number systems with digits 0 and 1. Our combinatorial approach is to construct infinitely many binary radix systems, each one from a single pair of binary strings. Every binary radix system that satisfies even a minimal set of conditions that would be expected of a positional number system can be constructed in this way.

1 Introduction

The terms *positional number system, radix system*, and *β-expansion* that appear in the literature all refer to the representation of real numbers in terms of a given base or radix B and a given finite set D of digits. Historically the base is 10 and the digit set is $\{0,1,2,\dots,9\}$ or, in the binary case, the base is 2 and the digit set is $\{0,1\}$. Alternative choices for the set of digits goes back at least to Cauchy, who suggested the use of negative digits in base 10, for example $D=\{-4,-3,\dots,4,5\}$. The *balanced ternary system* is a base 3 system with digit set $D=\{-1,0,1\}$ discussed by Knuth in [6]. In the balanced ternary, every integer, positive or negative, has a representation of the form $\sum_{n=0}^{N}\omega_n\,3^n$, where $\omega_n\in\{-1,0,1\}$ for all n. A well known system with digit set $\{0,1\}$ and base equal the golden ratio $(1+\sqrt{5})/2$ originated with [4]. Other early work on the representation of numbers using a non-integer base include those of Rényi [8] and Parry [7]. Positional number systems have also been extended to the representation of complex numbers [5] and, more generally, to the representation of points in $\mathbb{R}^d$ [10]. A vast array of additional references on positional number systems can be accessed by searching on the terms 'radix system' and 'β-expansion'.

Our intention in this paper is to provide a combinatorial framework for the binary representation of the real numbers. By binary we mean with digit set $\{0,1\}$. A precise definition of a *binary radix system* appears later in this section. Starting from any pair of infinite strings of $0's$ and $1's$ that satisfy very simple combinatorial conditions (given in Section 2), a binary radix system is constructed (in Section 3). Conversely, any binary radix system can be obtained by this combinatorial construction (Theorem 5.1 in Section 5). This leads to infinitely many binary radix systems, some of whose properties are investigated in this paper.

More specifically, our point of view is as follows. Let $\Omega = \{0,1\}^\infty$ denote the set of infinite strings of the form $\omega := \omega_0\,\omega_1\,\omega_2\cdots$, where $\omega_n \in \{0,1\}$ for all $n \geq 0$. Extend Ω to a set $\Omega^\bullet$ of "decimals" as follows.

$$\Omega^\bullet := \{\omega_0\,\omega_1\cdots\omega_N \,\bullet\, \omega_{N+1}\,\omega_{N+2}\cdots \,:\, \omega_0\,\omega_1,\,\omega_2\cdots \in \Omega\}\,.$$

In fact, we will refer to elements of $\Omega^\bullet$ as *decimals*. When no confusion arises, we omit an initial string of zeros before the decimal point and/or a terminal string of zeros after the decimal point, for example $01 \bullet 1000\cdots = 1 \bullet 1$. A line over a finite string denotes infinite repetition, for example $\overline{01} = 010101\cdots$.

If $\omega \in \Omega^\bullet$, then $\widehat{\omega} \in \Omega$ denotes the string ω with the decimal point removed. Moreover, if $\Gamma \subseteq \Omega^\bullet$, then $\widehat{\Gamma} = \{\widehat{\omega} \,:\, \omega \in \Gamma\}$. Conversely, if $\Gamma \subset \Omega$, then $\Gamma^\bullet$ denotes the set of all decimals obtained from the strings in Γ by placing the decimal point anywhere. For example if $\overline{10} \in \Gamma$, then $\{\dots, \bullet 00\overline{10},\, \bullet 0\overline{10},\, \bullet\overline{10},\, 1 \bullet \overline{01},\, 10 \bullet \overline{10}, \dots\} \subset \Gamma^\bullet$.

Definition 1.1. The **shift operator** $S : \Omega \to \Omega$ is defined by

$$S(\omega_0\,\omega_1\omega_2\cdots) = \omega_1\,\omega_2\cdots\,.$$

Let S^n denotes the n^{th} iterate of S. A subset $\Gamma \subset \Omega$ will be called **shift invariant** if $S^n(\Gamma) \subseteq \Gamma$. A subset $\Gamma \subset \Omega^\bullet$ will be called **shift invariant** if $[\,S^n(\widehat{\Gamma})\,]^\bullet = \Gamma$ for all $n \geq 0$. In other words, $\Gamma \subset \Omega^\bullet$ is shift invariant if, for any element of Γ, when we remove the decimal point, perform any number of shifts, and replace the decimal point anywhere, the result again lies in Γ.

The lexicographic order $\preceq$ on Ω is the total order defined by $\sigma \prec \omega$ if $\sigma \neq \omega$ and $\sigma_k < \omega_k$ where k is the least index such that $\sigma_k \neq \omega_k$. The lexicographic order on Ω can be extended to $\Omega^\bullet$ as follows. If

$$\begin{aligned} \sigma &= \sigma_0\,\sigma_1\cdots\sigma_N \,\bullet\, \sigma_{N+1}\,\sigma_{N+2}\cdots \quad \text{and} \\ \omega &= \omega_0\,\omega_1\cdots\omega_N \,\bullet\, \omega_{N+1}\,\omega_{N+2}\cdots \end{aligned}$$

(where some leading entries may possibly be 0's), then define $\sigma \prec \omega$ if $\sigma \neq \omega$ and $\widehat{\sigma} \prec \widehat{\omega}$ in the lexicographic order on Ω. For example $\bullet\overline{1} \prec 1 \bullet \overline{0}$ because $0 \bullet \overline{1} \prec 1 \bullet \overline{0}$ because $0\overline{1} \prec 1\overline{0}$.

The three conditions in the definition below are meant to be the minimum requirements that one would expect of a positional number system. Here $\mathbb{R}^+$ denote the set of non-negative real numbers.

Definition 1.2. A **binary radix system** is a triple (Γ, π, B), where $B > 1$ is a real number called the **base**, $\Gamma \subset \Omega^\bullet$ is called the **address space**, and $\pi : \Gamma \to \mathbb{R}^+$ is called the **radix map**. The following conditions must be satisfied:

1. Γ is shift invariant,
2. π is bijective and strictly increasing, and
3. for all $\omega_0\,\omega_1\cdots\omega_N \,\bullet\, \omega_{N+1}\,\omega_{N+2}\cdots \in \Gamma$ we have

$$\pi(\omega_0\,\omega_1\cdots\omega_N \,\bullet\, \omega_{N+1}\,\omega_{N+2}\cdots) = \sum_{n=-\infty}^{N} \omega_{N-n}\,B^n.$$

Conditions 2 and 3 are clear. Example 1.3 below should make condition 1 clear.

Example 1.3. The standard binary representation of the real numbers is an example, actually two examples. Let $B = 2$ and let Γ consist of all strings in $\Omega^\bullet$ except those ending in $0111\cdots$. Let $\pi(\omega_0\,\omega_1\cdots\omega_N\,\bullet\,\omega_{N+1}\,\omega_{N+2}\cdots) = \sum_{n=-\infty}^{N}\omega_{N-n}\,2^n$. Then (Γ,π,B) is a binary radix system. Alternatively, with the same $B = 2$ and radix map π, let Γ consist of all strings in $\Omega^\bullet$ except those ending in $1000\cdots$. Again (Γ,π,B) is a binary radix system. Note that an address space Γ with base $B = 2$ cannot contain both an element that ends in $1\overline{0}$ and an element that ends in $0\overline{1}$. This is because, by shift invariance, if this were so, then both $\bullet 1\overline{0}$ and $\bullet 0\overline{1}$ must lie in Γ. But $\pi(\bullet 1\overline{0}) = \pi(\bullet 0\overline{1})$, contradicting that π must be bijective. It is precisely to avoid this type of inconsistency, that some numbers are represented by a decimal ending in $1\overline{0}$ and others are represented by a decimal ending in $0\overline{1}$, that we require shift invariance as a condition for a binary radix system.

Example 1.4. In the introduction we referred to a well known radix system (Γ,π,B), where B is the golden ratio $(1+\sqrt{5})/2$. In this case, Γ consists of all strings in $\Omega^\bullet$ that do not contain 11 or $\overline{01}$ as a substring. Some elements in the address space are

$$0\bullet \qquad 1\bullet \qquad 10\bullet \qquad 100\bullet \qquad 101\bullet \qquad 1000\bullet \qquad 1001\bullet \qquad 1010\bullet \qquad 10000\bullet$$

The bijective map onto the positive reals is

$$\pi(\omega_0\,\omega_1\cdots\omega_N\,\bullet\,\omega_{N+1}\,\omega_{N+2}\cdots) = \sum_{n=-\infty}^{N}\omega_{N-n}\left(\frac{1+\sqrt{5}}{2}\right)^n.$$

That $\pi : \Gamma \to \mathbb{R}^+$ is bijective and strictly increasing is a special case of general results in Section 3.

The organization of the remainder of this paper is as follows. The notion of an *admissible* and *null* pair (α,β) of strings is introduced in Section 2. The construction of a binary radix system from an admissible non-null pair of strings is the subject of Section 3. Section 5 contains a proof that every binary radix system can be obtained by this construction from some admissible non-null pair (α,β). An algorithm is provided in Section 4 whose input is a particular binary radix system and a positive real number x and whose output is the decimal expansion of x in that binary radix system. For the standard base 2 radix system there is an associated tiling of the real line, where a tile consists of all those points on the real line with the same integer part. This is the trivial tiling consisting of unit length intervals. There is an an analogous tiling for any binary radix system but, in general, the tiles have various lengths, and the tiling is not periodic. This is the subject of Section 6.

2 Admissible and Null Strings

The notation

$$[\alpha,\beta] := \{\omega\in\Omega \,:\, \alpha\preceq\omega\preceq\beta\}$$

will be used for the closed interval; likewise for the open interval (α,β) and half open intervals $[\alpha,\beta)$ and $(\alpha,\beta]$.

Definition 2.1. Call a pair (α, β) of elements of Ω **admissible** if α and β satisfy the following two conditions.

1. $\alpha_0 = 0,\ \alpha_1 = 1 \quad$ and $\quad \beta_0 = 1,\ \beta_1 = 0,$
2. $S^n\alpha \notin (\alpha, \beta] \quad$ and $\quad S^n\beta \notin [\alpha, \beta) \quad$ for all $n \geq 0$.

Definition 2.2. The **address spaces** associated with such an admissible pair (α, β) are defined by

$$\begin{aligned}\Omega_{(\alpha,\beta,-)} &:= \{\omega \in \Omega \,:\, S^n\omega \notin (\alpha, \beta] \text{ for all } n \geq 0\},\\ \Omega_{(\alpha,\beta,+)} &:= \{\omega \in \Omega \,:\, S^n\omega \notin [\alpha, \beta) \text{ for all } n \geq 0\}.\end{aligned}$$

Moreover, let

$$\Omega(\alpha, \beta) := \Omega_{(\alpha,\beta,-)} \cup \Omega_{(\alpha,\beta,+)}.$$

Extend the definition of address spaces $\Omega_{(\alpha,\beta,-)}, \Omega_{(\alpha,\beta,+)}, \Omega_{(\alpha,\beta)}$ to address spaces $\Omega^\bullet_{(\alpha,\beta,-)}, \Omega^\bullet_{(\alpha,\beta,+)}, \Omega^\bullet_{(\alpha,\beta)}$ of decimals as follows:

$$\begin{aligned}\Omega^0_{(\alpha,\beta,-)} &:= \{\omega \in \Omega_{(\alpha,\beta,-)} \,:\, 0\omega \preceq \alpha\},\\ \Omega^0_{(\alpha,\beta,+)} &:= \{\omega \in \Omega_{(\alpha,\beta,+)} \,:\, 0\omega \prec \alpha\},\\ \Omega^0_{(\alpha,\beta)} &:= \Omega^0_{(\alpha,\beta,-)} \cup \Omega^0_{(\alpha,\beta,+)}.\end{aligned}$$

$$\begin{aligned}\Omega^\bullet_{(\alpha,\beta,-)} &:= [\,\Omega^0(\alpha, \beta, -)\,]^\bullet,\\ \Omega^\bullet_{(\alpha,\beta,+)} &:= [\,\Omega^0(\alpha, \beta, +)\,]^\bullet,\\ \Omega^\bullet_{(\alpha,\beta)} &:= [\,\Omega^0(\alpha, \beta)\,]^\bullet.\end{aligned}$$

The reason for introducing the spaces Ω^0 with the added condition $0\omega \prec \alpha$ is to insure that the spaces $\Omega^\bullet$ are shift invariant; see Proposition 2.7 below.

Example 2.3. If $\alpha = 0\overline{1} = 0\,1\,1\,1\cdots$ and $\beta = 1\overline{0} = 1\,0\,0\,0\cdots$, then (α, β) is an admissible pair. The address space $\Omega^\bullet_{(\alpha,\beta,-)}$ consists of all decimals that do not end in $1000\ldots$, and $\Omega^\bullet_{(\alpha,\beta,+)}$ consists of all decimals that do not end in $0111\ldots$. Notice that $\Omega^\bullet_{(\alpha,\beta,-)}$ and $\Omega^\bullet_{(\alpha,\beta,+)}$ are exactly the two address spaces for the standard binary radix system of Example 1.3.

Example 2.4. If $\alpha = \overline{01}$ and $\beta = 1\overline{0}$, then (α, β) is an admissible pair. Note that, although $11\overline{0} \in \Omega_{(\alpha,\beta)}$, the decimal $11\bullet \notin \Omega^\bullet_{(\alpha,\beta)}$. This is because $11\overline{0} \notin \Omega^0_{(\alpha,\beta)}$ since $011\overline{0} \succ \alpha$. Indeed, it is an easy exercise to show that $\omega \in \Omega^\bullet_{(\alpha,\beta)}$ if and only if ω does not contain 11 as a substring. Notice that $\Omega_{(\alpha,\beta,-)}$ is exactly the address space of the golden ratio based binary radix system of Example 1.4.

Definition 2.5. A string $\omega \in \Omega$ will be called **periodic** if $\omega = \overline{s}$ for some finite string s. A string $\omega \in \Omega$ will be called **eventually periodic** if there is an $m \geq 0$ such that $S^m\omega$ is periodic.

The fact that the address space $\Omega^\bullet_{(\alpha,\beta)}$ in Example 2.4 can be characterized by a set of "forbidden" finite substrings is not a coincidence, as explained in the proposition below.

Proposition 2.6. *If an admissible pair (α, β) is such that α and β are pereiodic, then there is a finite set T of finite strings such that $\Omega^{\bullet}_{(\alpha,\beta)}$ is the set of $\omega \in \Omega^{\bullet}$ such that s is not a substring of ω for any $s \in T$.*

More instructive than giving a formal proof of the proposition is to give an example that makes clear how a proof woould proceed. Referring to Definition 2.1 of admissible pair, if $\alpha = \overline{01101}$ and $\beta = \overline{100}$, then the forbidden set is $T = \{111, 11011, 000\}$. The string 111 is not allowed because $0111\gamma \succ \alpha$ for any string γ, and 11011 is not allowed because $011011\gamma \succ \alpha$ for any string γ. The string 000 is not allowed because $1000\gamma \prec \beta$ for any string γ. Because of the periodicity, no additional forbidden strings are required.

Note that the proposition is false if, instead of periodic, the hypothesis assumes only eventually periodic. For example, if $\alpha = 011\overline{01}$ and $\beta = 1\overline{0}$, then we would have to forbid an infinite set $111, 11011, 1101011, 110101011, \ldots$.

Proposition 2.7. *Given an admissible pair (α, β), the address spaces $\Omega^{\bullet}_{(\alpha,\beta,-)}$, $\Omega^{\bullet}_{(\alpha,\beta,+)}$, and $\Omega^{\bullet}_{(\alpha,\beta)}$ are shift invariant.*

Proof. The proposition follows in a straightforward way from the definition of the address spaces, except for one detail. Consider the case of $\Omega^{\bullet}_{(\alpha,\beta,-)}$ (the case $\Omega^{\bullet}_{(\alpha,\beta,+)}$ is similar). If $\omega := \omega_0 \cdots \omega_N \bullet \omega_{N+1}\omega_{N+2} \cdots \in \Omega^{\bullet}_{(\alpha,\beta,-)}$, then for $\Omega^{\bullet}_{(\alpha,\beta,-)}$ to be shift invariant it is necessary that $\bullet 0\, \omega_0\omega_1\omega_2 \cdots \in \Omega^{\bullet}_{(\alpha,\beta,-)}$. Therefore it is necessary, not only that $S^n\widehat{\omega} \notin (\alpha, \beta]$ for all $n \geq 0$, but also that $0\widehat{\omega} \notin (\alpha, \beta]$, i.e., $0\widehat{\omega} \preceq \alpha$. But this is exactly the condition in the defintion of $\Omega^{0}_{(\alpha,\beta,-)}$. $\square$

There exist admissible pairs (α, β) whose address spaces are exceptionally sparse in the following sense. For $\omega \in \Omega$ let $\omega|_n$ denote the initial finite segment $\omega_0\omega_1 \cdots \omega_{n-1}$ of length n. For $\Gamma \subseteq \Omega$, let $\Gamma_n := \{\gamma|_n : \gamma \in \Gamma\}$. Moreover $|\Gamma_n|$ denotes the cardinality of Γ_n. Define the **exponential growth rate** $h(\Gamma)$ of $\Gamma \subseteq \Omega$ by

$$h(\Gamma) = \limsup_{n\to\infty} \frac{1}{n} \ln |\Gamma_n|.$$

Consider Example 2.4. If $\alpha = 0\overline{1}$ and $\beta = 1\overline{0}$, then the corresponding address space $\Omega_{(\alpha,\beta)}$ for the standard binary radix system has largest possible exponential growth rate $h(\Omega_{(\alpha,\beta)}) = \ln(2)$ because $|(\Omega_{(\alpha,\beta)})_n| = 2^n$. Although the address spaces of admissible pairs usually have positive exponential growth rate, it is possible that $h(\Omega_{(\alpha,\beta)}) = 0$, as demonstated by the following example.

Example 2.8. The following α and β consititute an admissible pair:

$$\alpha = 01\overline{10}$$
$$\beta = 10\overline{01}.$$

First notice that a string $\omega \in \Omega_{(\alpha,\beta)}$ cannot cantain either 1000 or 0111 as a substring. Moreover ω cannot contain either $100\,s\,00$ or $011\,t\,11$ as a substring, where $s = 10101\cdots 1$ and $t = 01010\cdots 0$. In other words, ω is a string of alternating $0's$ and $1's$ with possibly one double zero or one double one, and possibly a string of zeros or a string of ones at the beginning. With these rigid restrictions, it is not hard to check that $h(\Omega_{(\alpha,\beta)}) = 0$.

Definition 2.9. If $h(\Omega_{(\alpha,\beta)}) = 0$, call the pair (α, β) **null**.

For an admissible pair (α, β) to be null is not typical. For example, if (α, β) is an admissible pair for which $\alpha = 011\alpha'$ and $\beta = 100\beta'$ and either α' begins wilth a 1 or β' begins with a 0, then (α, β) is non-null. This is because, in the case that α' begins wilth a 1 say, any string of the form $0(s_1)\,0\,(s_2)\,0\,(s_3)\,0\cdots \in \Omega_{(\alpha,\beta)}$, where $(s_n) = 1$ or $(s_n) = 11$ for $n = 1, 2, \dots$. It may be an interesting question to characterize the set of admissible pairs that are null.

3 The Radix Systems $R(\alpha, \beta, \pm)$

In the previous section, we constructed, for every admissible pair (α, β) of strings, two address spaces $\Omega^\bullet_{(\alpha,\beta,-)}$ and $\Omega^\bullet_{(\alpha,\beta,+)}$. In this section we describe how these two address spaces of decimals become the address spaces of two binary radix systems.

Theorem 3.1. *If (α, β) is an admissible non-null pair, then*

1. $\sum_{n=0}^\infty \alpha_n\, x^n = \sum_{n=0}^\infty \beta_n\, x^n$ *has a solution in the interval* $[\frac{1}{2}, 1)$ *of the real line, and*
2. *if b denotes the minimum such solution and $B_{(\alpha,\beta)} = 1/b$, then $(\Omega^\bullet_{(\alpha,\beta,-)}, \pi_{(\alpha,\beta)}, B_{(\alpha,\beta)})$ and $(\Omega^\bullet_{(\alpha,\beta,+)}, \pi_{(\alpha,\beta)}, B_{(\alpha,\beta)})$ are binary radix systems, where*

$$\pi_{(\alpha,\beta)}(\omega_0\,\omega_1 \cdots \omega_N \,\bullet\, \omega_{N+1}\,\omega_{N+2}\cdots) = \sum_{n=-\infty}^{N} \omega_{N-n}\, B^n_{(\alpha,\beta)}.$$

In the proof of Theorem 3.1 and also in the next section, the following notion is used.

Definition 3.2. Given an $x \in [0, 1)$, define the **projection map** $\pi_x : \Omega \to [0, 1]$ by

$$\pi_x(\omega) := (1 - x)\sum_{k=0}^{\infty} \omega_k\, x^k.$$

Note that $\pi_x(\overline{0}) = 0$ and $\pi_x(\overline{1}) = 1$ for all $x \in [0, 1)$.

Proof. (Theorem 3.1) Statement 1 is in [2, Theorem 13], a main result in that paper. That (α, β) is admissible and non-null is crucial for that result. It is also proved in that paper that $\pi_b : \Omega_{(\alpha,\beta,\pm)} \to [0, 1]$ is strictly increasing and bijective. The shift invariance of $\Omega^\bullet_{(\alpha,\beta,\pm)}$, which is condition 1 in Definition 1.2, is Lemma 2.7.

It now suffices to prove that $\pi_{(\alpha,\beta)} : \Omega^\bullet_{(\alpha,\beta,\pm)} \to \mathbb{R}^+$ is strictly increasing and bijective. We will prove it for $\Omega^\bullet_{(\alpha,\beta,+)}$. The proof for $\Omega^\bullet_{(\alpha,\beta,-)}$ is essentially the same. Abbreviate $B := B_{(\alpha,\beta,\pm)}$. Referring to statement 1 in the theorem, let $p = \sum_{n=0}^\infty \alpha_n\, b^n = \sum_{n=0}^\infty \beta_n\, b^n$. Assume that $\omega \in \Omega_{(\alpha,\beta,+)}$. Since $\pi_b : \Omega_{(\alpha,\beta,+)} \to [0, 1]$ is strictly increasing, we have $0\,\omega \prec \alpha$ if and only if $b\pi_b(\omega) = \pi_b(0\omega) < \pi_b(\alpha) = p$ if and only if $\pi_b(\omega) < Bp$. Therefore $\pi_b : \Omega^0_{(\alpha,\beta,+)} \to [0, Bp]$ is strictly increasing and bijective. Now let $\Omega^\bullet_N$ denote the set of elements of $\Omega^\bullet_{(\alpha,\beta,+)}$ of the form $\omega := \omega_0\omega_1\omega_2\cdots\omega_{N-1} \bullet \omega_N\omega_{N+1}\cdots$, where $\widehat{\omega} := \omega_0\omega_1\omega_2\cdots \in \Omega^0_{(\alpha,\beta,+)}$. The relationship between the projection map π_b and the radix map $\pi_{(\alpha,\beta)}$ is given by $\pi_{(\alpha,\beta)}(\omega) = \frac{B^N}{1-b}\,\pi_b(\widehat{\omega})$, and therefore $\pi_{(\alpha,\beta)} : \Omega^\bullet_N \to [\,0, \frac{B^{N+1}p}{b-1}\,]$ is a bijection and strictly increasing. Since the sequence $\Omega^\bullet_0 \subset \Omega^\bullet_1 \subset \Omega^\bullet_2 \subset \cdots$ of sets is nested, and since $\bigcup_{N=0}^\infty \Omega^\bullet_N = \Omega^\bullet_{(\alpha,\beta,+)}$, the proof is complete. □

Definition 3.3. A two radix systems $(\Omega^\bullet_{(\alpha,\beta,\pm)}, \pi_{(\alpha,\beta)}, B_{(\alpha,\beta)})$ constructed from an admissible non-null pair (α, β) as in Theorem 3.1 will be called (α, β)-**radix systems** and denoted $R(\alpha, \beta, \pm)$.

Example 3.4. Continuing from Example 2.3, consider the binary radix systems $R(0\overline{1}, 1\overline{0}, \pm)$. The number b in Theorem 3.1 is the smallest solution to the equation $\sum_{n=1}^{\infty} x^n = 1$, which, because the right hand side is a geometric series, reduces to $2x = 1$. So $b = 1/2$ and the base is $B_{(\alpha,\beta)} = 1/b = 2$. The radix map is

$$\pi_{(\alpha,\beta)}(\omega_0\,\omega_1 \cdots \omega_N \bullet \omega_{N+1}\,\omega_{N+2} \cdots) = \sum_{n=-(N)}^{\infty} \omega_{n+N} \left(\frac{1}{2}\right)^n = \sum_{n=-\infty}^{N} \omega_{N-n}\, 2^n.$$

The address space $\Omega^\bullet_{(\alpha,\beta,-)}$ consists of all decimal strings that do not end in $1000\cdots$, and $\Omega^\bullet_{(\alpha,\beta,+)}$ consists of all decimal strings that do not end in $0111\cdots$. This example is the standard binary representaion of of the non-negative real numbers given in Example 1.3.

Example 3.5. The two binary radix systems $R(\overline{01}, 1\overline{0}, \pm)$ are a continuation of Example 2.4, in which $\alpha = \overline{01}$ and $\beta = 1\overline{0}$ form an admissible non-null pair. The number b in Theorem 3.1 is the smallest solution to the equation $\sum_{n=0}^{\infty} x^{2n+1} = 1$, which reduces to $x^2 + x - 1 = 0$. So $b = (\sqrt{5} - 1)/2$, and the base $B_{(\alpha,\beta)} = 1/b = (1 + \sqrt{5})/2$ is the golden ratio. The radix map is

$$\pi_{(\alpha,\beta)}(\omega_0\,\omega_1 \cdots \omega_N \bullet \omega_{N+1}\,\omega_{N+2} \cdots) = \sum_{n=-\infty}^{N} \omega_{N-n} \left(\frac{1+\sqrt{5}}{2}\right)^n.$$

This is the golden ratio based representation of the reals introduced in Example 1.4.

Example 3.6. Consider the(α, β)-radix systems $R(\alpha_1, \beta_1, \pm)$, $R(\alpha_2, \beta_2, \pm)$, and $R(\alpha_3, \beta_3, \pm)$, where the admissible non-null pairs are:

$$\begin{array}{lll} \alpha_1 = \overline{01000} & \alpha_2 = \overline{011} & \alpha_3 = \overline{01} \\ \beta_1 = 1\overline{0} & \beta_2 = \overline{10} & \beta_3 = \overline{100}. \end{array}$$

In all three cases the number b in Theorem 3.1 is the solution to the equation $x^3 + x^2 - 1 = 0$ in the interval $[0, 1)$; approximately $b \approx 0.7549$. Therefore the base $B_{(\alpha,\beta)} \approx 1.3247$ and the radix maps $\pi_{(\alpha,\beta)}$ are the same in all three cases. The three address spaces, however, are pairwise distinct. The address space $\Omega^\bullet_{(\alpha_1,\beta_1)}$ consists of all decimals that do not contain the substrings $11, 101, 1001$ or 10001. The address space $\Omega^\bullet_{(\alpha_2,\beta_2)}$ consists of all decimals that do not contain the substrings 100 or 111. The address space $\Omega^\bullet_{(\alpha_3,\beta_3)}$ consists of all decimals that do not contain the substrings 11 or 1000.

4 Algorithm for Determining the Address

Given a binary radix system, the radix map assigns a non-negative real number to each decimal in the address space. In this section an algorithm is provided for converting in the opposite direction. Given a non-negative real number x, the algorithm determines its decimal representation in the binary radix system. More precisely, if $R(\alpha, \beta, \pm) =$

$(\Omega^{\bullet}_{(\alpha,\beta,\pm)}, \pi_{(\alpha,\beta)}, B_{(\alpha,\beta)})$ and $x \in \mathbb{R}^{+}$, then the algorithm finds $\sigma \in \Omega^{\bullet}_{(\alpha,\beta,-)}$ and $\omega \in \Omega^{\bullet}_{(\alpha,\beta,+)}$ such that $\pi_{(\alpha,\beta)}(\sigma) = \pi_{(\alpha,\beta)}(\omega) = x$.

We begin by defining a certain family of functions and introduce notation for the itineraries of points for this family of functions. Given B such that $1 < B \leq 2$ and p such that $1 - b \leq p \leq b$ where $b = 1/B$, consider the two functions $f_{(B,p,\pm)} : [0,1] \to [0,1]$ defined by

$$f_{(B,p,-)}(x) := \begin{cases} Bx & \text{if } 0 \leq x \leq p \\ Bx + (1-B) & \text{if } x > p. \end{cases} \tag{1}$$

and

$$f_{(B,p,+)}(x) := \begin{cases} Bx & \text{if } 0 \leq x < p \\ Bx + (1-B) & \text{if } x \geq p, \end{cases} \tag{2}$$

The facts that $1 < B \leq 2$ and $1 - b \leq p \leq b$ guarantee that $f_{(\alpha,\beta,\pm)}$ have the form shown in Figure 1. For a function f, the n^{th} iterate, i.e. f composed with itself n times, is denoted f^n.



Figure 1: The function $f_{(B,p,\pm)}$.

Definition 4.1. Define the two **itinerary maps** $\widehat{\tau}_{(B,p,\pm)} : [0,1] \to \Omega$ by

$$[\widehat{\tau}_{(B,p,-)}(y)]_k = \begin{cases} 0 & \text{if } f^k_{(B,p,-)}(y) \leq p \\ 1 & \text{if } f^k_{(B,p,-)}(y) > p, \end{cases}$$

and

$$[\widehat{\tau}_{(B,p,+)}(y)]_k = \begin{cases} 0 & \text{if } f^k_{(B,p,+)}(y) < p \\ 1 & \text{if } f^k_{(B,p,+)}(y) \geq p. \end{cases}$$

In dynamical systems terminology, $\widehat{\tau}_{(\alpha,\beta,-)}(y)$ and $\widehat{\tau}_{(\alpha,\beta,+)}(y)$ are called the **itineraries** of the point y.

Algorithm

Input: An admissible and non-null pair of strings (α, β) and an $x \in \mathbb{R}^+$.

Output: The decimals representations σ and ω for x in the binary radix systems $R(\alpha, \beta, -)$ and $R(\alpha, \beta, -)$, respectively.

1. Find the least $x \in (0,1)$ such that $\sum_{n=0}^{\infty} \alpha_n\, x^n = \sum_{n=0}^{\infty} \beta_n\, x^n$; call it b.
2. Compute $p := \pi_b(\alpha) = \pi_b(\beta)$, where π_b is the projection map.
3. Find the minimum non-negative integer N such that $y := b^N(1-b)x < p$.
4. Compute $\widehat{\tau}_{(\alpha,\beta,-)}(y) = \sigma_0\,\sigma_1\,\sigma_2 \cdots$ and $\widehat{\tau}_{(\alpha,\beta,+)}(y) = \omega_0\,\omega_1\,\omega_2 \cdots$.

4. Return

$$\sigma = \sigma_0\,\sigma_1\,\sigma_2 \cdots \sigma_N \bullet \sigma_{N+1}\,\sigma_{N+2} \cdots , \quad \text{and}$$
$$\omega = \omega_0\,\omega_1\,\omega_2 \cdots \omega_N \bullet \omega_{N+1}\,\omega_{N+2} \cdots .$$

Denote the output σ and ω of the algorithm by $\tau_{(\alpha,\beta,-)}(x)$ and $\tau_{(\alpha,\beta,+)}(x)$, respectively. The following theorem suffices to prove the validity of the algorithm.

Theorem 4.2. *The map* $\tau_{(\alpha,\beta,\pm)} : \mathbb{R}^+ \to \Omega^{\bullet}_{(\alpha,\beta,\pm)}$ *is the inverse of* $\pi_{(\alpha,\beta)} : \Omega^{\bullet}_{(\alpha,\beta,\pm)} \to \mathbb{R}^+$.

Proof. We will prove that $\tau_{(\alpha,\beta,-)}$ is the inverse of $\pi_{(\alpha,\beta)}$; a similar proof holds for $\tau_{(\alpha,\beta,+)}$. We first show that the functions $f_{(B,p,\pm)}$ are well defined, i.e. $1 < B \le 2$ and $1 - b \le p \le b$. Clearly $1 < B \le 2$ because $1 > b \ge 1/2$ as stated in Theorem 3.1. Moreover $1 - b \le p \le b$ because

$$1 - b \le (1-b)\sum_{n=0}^{\infty} \beta_n b^n = p = (1-a)\sum_{n=0}^{\infty} \alpha_n b^n \le (1-a)\sum_{n=1}^{\infty} b^n = b.$$

We next show that $\pi_b \circ \widehat{\tau}_{(B,p,-)}$ is the identity on $[0,1]$. If $f_0(x) = Bx$, $f_1(x) = Bx + (1-B)$, and $g_0(x) = bx$, $g_1(x) = bx + (1-b)$, then f_0 and g_0 are inverses, as are f_1 and g_1. Expressing $g_i(x) = bx + i(1-b)$ for $i = 0, 1$ and iterating

$$g_{\omega_0} \circ g_{\omega_1} \circ g_{\omega_2} \circ \cdots \circ g_{\omega_k}(x) = b^k x + (b^{k-1}\omega_{k-1} + \cdots + a\omega_1 + \omega_0)(1-b).$$

Therefore for any x_0 we have

$$\lim_{k\to\infty} g_{\omega_0} \circ g_{\omega_1} \circ g_{\omega_2} \circ \cdots \circ g_{\omega_k}(x_0) = (1-b)\sum_{k=0}^{\infty} \omega_k\, b^k = \pi_b(\omega).$$

Let $M_0 = [0,p]$ and $M_1 = (p,1]$. Let $x \in [0,1]$ and $\widehat{\tau}_{(B,p,-)}(x) = \omega$. It follows from the definition of $\widehat{\tau}_{(B,p,-)}$ that

$$x \in M_{\omega_0}, \quad f_{\omega_0}(x) \in M_{\omega_1}, \quad f_{\omega_1} \circ f_{\omega_0}(x) \in M_{\omega_2}, \quad f_{\omega_2} \circ f_{\omega_1} \circ f_{\omega_0} \in M_{\omega_3}, \ldots$$

and therefore

$$x \in g_{\omega_0}(M_{\omega_1}), \quad x \in g_{\omega_0} \circ g_{\omega_1}(M_{\omega_2}), \quad g_{\omega_0} \circ g_{\omega_1} \circ g_{\omega_2}(M_{\omega_3}), \ldots .$$

Hence $(\pi_b \circ \widehat{\tau}_{(\alpha,\beta,-)})(x) = \lim_{k\to\infty} g_{\omega_0} \circ g_{\omega_1} \circ g_{\omega_2} \circ \cdots \circ g_{\omega_k}(x_0) = x$.

To show that $\tau_{(\alpha,\beta,-)}$ is the inverse of $\pi_{(\alpha,\beta)}$, with notation as in the algorithm, abbreviating $\tau_{(B,p,-)}$ to τ, and letting $\tau(x) = \sigma$ we have

$$(\pi_{(\alpha,\beta)} \circ \tau)(x) = \pi_{(\alpha,\beta)}(\sigma) = \frac{B^N}{1-b}\,\pi_b(\widehat{\sigma}) = \frac{B^N}{1-b}\,(\pi_b \circ \widehat{\tau})(y) = \frac{B^N}{1-b}\,b^n(1-b)x = x.$$

It remains to show that the image of any $x \in \mathbb{R}^+$ under the map $\tau_{(\alpha,\beta,-)}$ lies in $\Omega^{\bullet}_{(\alpha,\beta,-)}$. With y as defined in the algorithm, it is sufficient to show that $\widehat{\tau}_{(\alpha,\beta,-)}(y) \in \Omega^{0}_{(\alpha,\beta,-)}$. Let τ_- and τ_+ denote the itineraries of the point p of the functions $f_{(B,p,-)}$ and $f_{(B,p,+)}$, respectively. In [1, Theorem 5.1] it is proved that $\widehat{\tau}_{(\alpha,\beta,-)}([0,1]) = \{\omega \in \Omega \,:\, S^n\omega \notin (\tau_-,\tau_+] \text{ for all } n \geq 0\}$, and in [2, Theorem 1] it is proved that $\alpha = \tau_-$ and $\beta = \tau_+$. Therefore $\widehat{\tau}_{(B,p,-)}([0,1]) = \Omega_{(\alpha,\beta,-)}$. It only remains to show that if $\widehat{\omega} = \widehat{\tau}_{(B,p,-)}(y)$, then $0\widehat{\omega} \preceq \alpha$. However, since $y < p$ as in step 1 of the algorithm, it follows immediatly from the definition of the itinerary $\widehat{\tau}_{(B,p,-)}(y)$ of y that $\widehat{\omega}_0 = 0$. In particular $0\widehat{\omega} \prec \alpha$. ☐

Corollary 1. *For each base B, $1 < B < 2$, there exist infinitely many binary radix systems with base B.*

Proof. Given B, let p be any real number such that $1 - 1/B \leq p \leq 1/B$, and consider the two functions f_- and f_+ defined exactly as in Equations (1) and (2), respectively. If α_p is the itinerary of the point p for the function f_-, and β_p is the itinerary of the function f_+, then (α_p, β_p) is an admissible non-null pair. The proof of this fact appears in [1, Theorem 5.2]. Therefore $R(\alpha_p,\beta_p,-)$ and $R(\alpha_p,\beta_p,-)$ are binary radix systems. For each B, however, there are infintely many choices for p. Each choice of p leads to a distinct addmissible pair (α_p,β_p) because, according to [9, Lemma 3.1], α_p and β_p are strictly increasing as functions of p. ☐

5 Every Binary Radix System is an (α,β)-Radix System

This section is dedicated to the proof of the following theorem.

Theorem 5.1. *For every binary radix system (Γ,π,B), there is an admissible non-null pair (α,β) such that either $R(\alpha,\beta,-) = (\Gamma,\pi,B)$ or $R(\alpha,\beta,+) = (\Gamma,\pi,B)$.*

Proof. Let

$$\alpha = \sup\{\gamma \in \widehat{\Gamma} \,:\, \gamma_0 = 0\},$$
$$\beta = \inf\{\gamma \in \widehat{\Gamma} \,:\, \gamma_0 = 1\}.$$

It follows readily from from these definitions and the shift invariance of Γ that

$$S^n\alpha \notin (\alpha,\beta) \quad \text{and} \quad S^n\beta \notin (\alpha,\beta)$$

for all $n \geq 0$, which is close to, but not quite, the definition of an admissible pair. Condition 1 in Definition 2.1 of admissible pair holds because, by the shift invariance of Γ, there are strings in Γ_0 that begin with 01 and stings in Γ_1 that begin with 10.

To show that (α,β) is an admissible pair, it only remains to prove that there is no $n \geq 0$ such that $S^n\alpha = \beta$ and no $n \geq 0$ such that $S^n\beta = \alpha$. The fact that there is no element of $\widehat{\Gamma}$ between α and β in the lexicographic order and that π is increasing and surjective forces $\pi(\bullet\alpha) = \pi(\bullet\beta)$. Moreover, either $\bullet\alpha \in \Gamma$ or $\bullet\beta \in \Gamma$, but not both. We will assume that $\bullet\alpha \in \Gamma$ and $\bullet\beta \notin \Gamma$; the proof in the case that $\bullet\beta \in \Gamma$ is essentially the same. There is no $n \geq 0$ such that $S^n\alpha = \beta$; otherwise the fact that $\alpha \in \widehat{\Gamma}$ and the shift invariance of Γ (and hence the shift invariance of $\widehat{\Gamma}$) would imply that $\beta \in \widehat{\Gamma}$, a contradiction. Finally assume, by way of contradiction, that there is an $n \geq 0$ such that $S^n\beta = \alpha$. Then $\beta = t\alpha \prec t\beta$, where t is a finite string. If there exists a $\gamma \in \widehat{\Gamma}$ such that $t\alpha = \beta \prec \gamma \prec t\beta$, then $\alpha = S^n(t\alpha) \prec S^n\gamma \prec S^n(t\beta) = \beta$. But by the definition of α and β as sup and inf, there can be no such $\gamma \in \widehat{\Gamma}$. Therefore there is no such γ with $\beta \prec \gamma \prec t\beta$, which contradicts the definition of β as $\inf\{\gamma \in \widehat{\Gamma} : \gamma_0 = 1\}$.

The claim is now that $(\Gamma, \pi, B) = R(\alpha, \beta, -) := (\Omega^{\bullet}_{(\alpha,\beta,-)}, \pi_{(\alpha,\beta)}, B_{(\alpha,\beta)})$. To prove this, and thus finish the proof, it must be shown that

1. $\Gamma = \Omega^{\bullet}_{(\alpha,\beta,-)}$,
2. $B = B_{(\alpha,\beta)}$, and
3. (α, β) is not null.

Concerning item 1, the definition of α and β implies that $\widehat{\Gamma} \subseteq \Omega_{(\alpha,\beta,-)}$. That Γ is shift invariant further implies that $\widehat{\Gamma} \subseteq \Omega^{0}_{(\alpha,\beta,-)}$, and therefore that $\Gamma \subseteq \Omega^{\bullet}_{(\alpha,\beta,-)}$. To prove equality we take a somewhat circuitious route.

Define

$$\Gamma' = \{t\gamma \,:\, \gamma \in \widehat{\Gamma},\, t \,\text{a finite string of}\, 1's \,\text{including the empty string}\}.$$

For $b = 1/B$, the projection map $\pi_b : \Gamma' \to [0,1]$ as in Definition 3.2 is strictly increasing and surjective. To see this, first note that, since π is strictly increasing on Γ, the map $\pi_b : \widehat{\Gamma} \to [0,q]$ is stringly increasing and surjective for some $0 < q \leq 1$. Moreover, if $\omega \in \Gamma' \setminus \widehat{\Gamma}$ and $\gamma \in \widehat{\Gamma}$, then, by Definition 2.2 of the address spaces (recall that $\Gamma \subseteq \Omega^{\bullet}_{(\alpha,\beta,-)}$), we have $0\gamma \preceq \alpha \prec 0\omega$. Therefore $\gamma \prec \omega$, i.e., every element of $\widehat{\Gamma}$ is less than every element of $\Gamma' \setminus \widehat{\Gamma}$. In addition, since $0\gamma \preceq \alpha$, we have $\pi_b(\gamma) \leq (1/b)\,\pi_b(\alpha)$ by the fact that π is stringly increasing on Γ. And similarly, since $\alpha \prec 0\omega$, we have $(1/b)\,\pi_b(\alpha) < \omega$. Therefore $\pi_b(\gamma) < \pi_b(\omega)$. Finally, if σ and ω both lie in $\Gamma' \setminus \widehat{\Gamma}$ and $\sigma \prec \omega$, we will show that $\pi_b(\sigma) < \pi_b(\omega)$. Let $\sigma = t_1\gamma_1 \in \Gamma' \setminus \widehat{\Gamma}$, where t_1 is a string of m ones and $\gamma_1 \in \widehat{\Gamma}$, and $\omega = t_2\gamma_2 \in \Gamma' \setminus \widehat{\Gamma}$, where t_2 is a string of $n \geq m$ ones and $\gamma_2 \in \widehat{\Gamma}$. Then $t\gamma_2 \succ \gamma_1$, where t is a string of $n - m$ ones. Therefore $\pi_b(t\gamma_2) \succ \pi_b(\gamma_1)$ and hence $\pi_b(\omega) \succ \pi_b(\sigma)$. Thus π_b is strictly increasing on Γ'.

To show that $\pi_b : \Gamma' \to [0,1]$ is surjective, we express Γ' as the union of non-overlapping intervals and show that the images of these intervals under π_b leave no gaps. First note that the greatest element of $\widehat{\Gamma}$ is $S\alpha$; the smallest element of $1\widehat{\Gamma} \setminus \widehat{\Gamma}$ is $1S^2\alpha$ and the largest is $1S\alpha$; the smallest element of $11\widehat{\Gamma} \setminus (\widehat{\Gamma} \cup 1\widehat{\Gamma})$ is $11S^2\alpha$ and the largest is $11S\alpha$; etc. However, $\pi_b(1S^2\alpha) = \pi_b(S\alpha)$, and therefore $\pi_b(11S^2\alpha) = \pi_b(1S\alpha)$, etc. Hence $\pi_b : \Gamma' \to [0,1]$ is surjective.

A map $\tau : [0,1] \to \Omega$ such that $\pi_b \circ \tau$ is the identify is called a *section* of $\pi_b : \Omega \to [0,1]$. If $\tau([0,1])$ is shift invariant, then τ is called a *shift invariant section*. In our case, the inverse of π_b restricted to Γ' is a shift invariant section. Call this section τ_b. A *mask* (in our case) is a partition of the interval $[0,1]$ into two parts M_0 and M_1. Given a mask $M = \{M_0, M_1\}$, define a function

$$f_{(B,M)} = \begin{cases} Bx & \text{if } x \in M_0 \\ BX + (1-B) & \text{if } x \in M_1. \end{cases}$$

The *itinerary map* $\tau_{(B,M)} : [0,1] \to \Omega$ is defined by

$$[\tau_{(B,M}(x)]_k = \begin{cases} 0 & \text{if } f^k_{(B,M)}(x) \in M_0 \\ 1 & \text{if } \; f^k_{(B,M)}(x) \in M_1. \end{cases}$$

To continue the proof, we use [3, Theorem 4] which states the following. The itinerary map $\tau_{(B,M)}$ is a section of π_b, and conversely, every shift invariant section of π_b is of the above form. Hence $\tau = \tau_{(B,M)}$ for some mask M. Because τ is increasing and because $0\sigma \prec 1\omega$ for any $\sigma, \omega \in \Omega$, the mask must be of the form $M_0 = [0,p]$, $M_1 = (p,1]$ for some $p \in (0,1)$. In particular, $\tau_b = \widehat{\tau}_{(B,p,-)}$ as in Definition 4.1 for some $p \in (0,1)$. Let τ_- and τ_+ denote the itineraries $\widehat{\tau}_{(B,p,-)}(p)$ and $\widehat{\tau}_{(B,p,+)}(p)$, respectively, of the point p. Then $\Omega_{(\tau_-,\tau_+,-)} = \tau([0,1]) = \Gamma' \subseteq \Omega_{(\alpha,\beta,-)}$, the first equality by [1, Theorem 5.1]. If $\Gamma' \neq \Omega_{(\alpha,\beta,-)}$, then either $\tau_- \prec \alpha$ or $\tau_+ \succ \beta$. In either case there is a contradiction to the definition of α as a sup or β as an inf. Therefore $\Gamma' = \Omega_{(\alpha,\beta,-)}$, which implies that $\widehat{\Gamma} = \Omega^0_{(\alpha,\beta,-)}$, which in turn implies that $\Gamma = \Omega^\bullet_{(\alpha,\beta,-)}$.

To prove items 2 and 3 in the list, we use a result of Parry [7, Page 373] on the topological entropy of a dynamical system on the unit interval, where the function is of the form $f_{(B,p,\pm)}$ shown in Figure 1. In particular it provides the first equaliy in

$$\ln(B) = h(\Omega_{(\tau_-,\tau_+,-)}) = h(\Omega_{(\alpha,\beta,-)}) = \ln(B_{(\alpha,\beta)}).$$

This proves both that $B = B_{(\alpha,\beta)}$, and also that (α,β) is non-null since $B > 1$ in a binary radix system implies that $h(\Omega_{(\alpha,\beta,-)}) > 0$. $\square$

6 Radix Tilings of the Real Line

For an element $\omega \in \Omega^\bullet$, let $\omega\bullet$ denote the finite substring of ω to the left of the decimal point.

Definition 6.1. Given a binary radix system (Γ, π, B) and a finite string s, let

$$T'_s := \{\pi(\omega) \, : \, \omega \in \Gamma \;\text{ and }\; \omega\bullet = s\}.$$

Note that, for many values of s, the set T'_s may be empty. For instance, in Example 1.4 the set $T'_{011} = \emptyset$. If $T'_s \neq \emptyset$, then the closure T_s of T'_s is a closed interval which we call a **tile**. If Ω_F denotes the set of all finite strings, let

$$\mathcal{T} := \{T_s \, : \, s \in \Omega_F\}.$$

Then $\mathcal{T}$ is a collection of non-overlapping intervals whose union is $\mathbb{R}^+$. The set $\mathcal{T}$ will be referred to as the **tiling** of $\mathbb{R}^+$ for the binary radix system. For an (α,β)-radix system, the corresponding tiling is denoted by $\mathcal{T}_{(\alpha,\beta)}$. The tiling for $R(\alpha,\beta,-)$ is the same as for $R(\alpha,\beta,-)$.

Example 6.2. For the standard binary radix system (Examples 1.3, 2.3, and 3.4), the tiling is the set $\mathcal{T} = \{[n, n+1] \ : \ n \geq 0\}$ of unit length intervals.

Example 6.3. For the golden ratio based radix system (Examples 1.4, 2.4, and 3.5), there are tiles of two lengths in the ratio $1 : \frac{1+\sqrt{5}}{2}$. The tiling $\mathcal{T}$ is a well known non-periodic tiling of the line. If the tiles are denoted 1 and B (for their lengths), then the sequence of tiles in the tiling of $\mathbb{R}^+$, from left to right, is

$$B\,1\,B\,B\,1\,B\,1\,B\,B\,1\,B\,B\,1\,B\,1\,B\,B\,1\,B\,1\,B\,B\,1\cdots. \tag{1}$$

This tiling is self-replicating in the following sense. If $f_B : \mathbb{R}^+ \to \mathbb{R}^+$ is the function $f_B(x) = Bx$, then $B(\mathcal{T}) := \{f_B(T) : T \in \mathcal{T}\}$ is also a tiling of $\mathbb{R}^+$, and each tile in $B(\mathcal{T})$ is the union of tiles in $\mathcal{T}$. Moreover, the sequence (1) above can be recursively generated, starting from B and using the substitution rules

$$\begin{aligned} B &\leftarrow B1 \\ 1 &\leftarrow B. \end{aligned}$$

In other words, the tiling is recursively generated as follows:

$$B \to B1 \to B1B \to B1BB1 \to B1BB1B1B \to \cdots.$$

Definition 6.4. With notation as in the example above, call a tiling $\mathcal{T}$ **self-replicating** if each tile in $B(\mathcal{T})$ is the union of tiles in $\mathcal{T}$.

Theorem 6.5. *Let (α, β) be an admissible pair.*

1. *The tiling $\mathcal{T}_{(\alpha,\beta)}$ is self-replicating.*
2. *If α and β are eventually periodic, then there are at most finitely many lengths of tiles in the tiling $\mathcal{T}_{(\alpha,\beta)}$.*

Proof. Let (Γ, π, B) be the binary radix system. Concerning statement 1, consider the set $X = \{x_1, x_2, \dots\}$ of right endpoints of the tiles in $\mathcal{T}_{(\alpha,\beta)}$. It suffices to prove that $Bx_n \in X$ for all $n \geq 1$. For any $n \geq 1$, the string $\tau_{(\alpha,\beta,-)}(x_n)$ must be of the form $\sigma = s\alpha_0\alpha_1\cdots\alpha_N \bullet \alpha_{N+1}\cdots$ for some finite string s and some $N \geq 0$. This is because, if $\sigma' = s\beta_0\beta_1\cdots\beta_N \bullet \beta_{N+1}\cdots$, then $\pi(\sigma) = \pi(\sigma')$ is the endpoint of two consecutive intervals in $\mathcal{T}_{(\alpha,\beta)}$. By the shift invariance of the address space, $\omega := s\alpha_0\alpha_1\cdots\alpha_N\alpha_{N+1}\bullet \alpha_{N+2}\cdots$ also lies in $\Omega^\bullet_{(\alpha,\beta,-)}$ and and hence, ω being of the above form, $\pi_{(\alpha,\beta)}(\omega)$ also lies in X. Moreover $Bx_n = B\pi_{(\alpha,\beta)}(\sigma) = \pi_{(\alpha,\beta)}(\omega) \in X$.

Concerning statement 2, from the form of the decimal representation of the endpoints of the intervals of the tiling (as discussed above), the lengths of the tiles are of the form $\pi_{(\alpha,\beta)}(\bullet S^m\alpha) - \pi_{(\alpha,\beta)}(\bullet S^n\beta)$ for some $n, m \geq 0$. Since α and β are eventually periodic, there are at most finitely many shifted strings $S^m\alpha$ and $S^n\beta$. $\square$

Example 6.6. This is a continuation of Example 3.6 in which three distinct (α, β)-radix systems have the same base B. The three associated self-replicating tilings are also distinct. From the self-replicating property, it is routine to find, in each of the three cases, substitution rules for the tiling.

Referring to the notation in Example 3.6, for the tiling $\mathcal{T}_{(\alpha_1,\beta_1)}$ there are five tile types $T_0, T_1, T_2, T_3, T_4, T_5$ of relative lengths $1, B, B^2, B^3, B^4$, respectively. To simplify notation we refer to tile T_i as simply i. Then the substitution rules are:

$$0 \leftarrow 1 \qquad 1 \leftarrow 2 \qquad 2 \leftarrow 3 \qquad 3 \leftarrow 4 \qquad 4 \leftarrow 40,$$

and the tiling begins, from left to right:

$$4\,0\,1\,2\,3\,4\,4\,0\,4\,0\,1\,4\,0\,1\,2\,4\,0\,1\,2\,3\,4\;\;0\,1\,2\,3\,4\,4\,0\,1\,2\,3\,4\,4\,0\ldots.$$

For the tiling $\mathcal{T}_{(\alpha_2,\beta_2)}$ there are 4 tile types T_0, T_1, T_2, T_3, T_4 of relative lengths $1, B, B^2, B^3, B^6 = (B+1)^2$, respectively. The substitution rules are:

$$0 \leftarrow 1 \qquad 1 \leftarrow 2 \qquad 2 \leftarrow 10 \qquad 3 \leftarrow 32,$$

and the tiling begins, from left to right:

$$3\,2\,1\,0\,2\,1\,1\,0\,2\,2\,1\,1\,0\,1\,0\,2\,2\,1\,2\,1\,1\;\;0\,1\,0\,2\ldots.$$

For the tiling $\mathcal{T}_{(\alpha_3,\beta_3)}$ there are 4 tile types T_0, T_1, T_2, T_3, T_4 of relative lengths $1, B, B^2, B^3, 1 + B + B^2$, respectively. The substitution rules are:

$$0 \leftarrow 1 \qquad 1 \leftarrow 2 \qquad 2 \leftarrow 01 \qquad 3 \leftarrow 31,$$

and the tiling begins, from left to right:

$$3\,1\,2\,0\,1\,1\,2\,2\,0\,1\,0\,1\,1\,2\,1\,2\,2\,0\,1\,2\,0\;\;1\,0\,1\,1\,2\ldots.$$

All three of the above tilings are self-replicating under expansion by the factor B that is the real root of the equation $B^3 - B - 1 = 0$.

References

[1] M. Barnsley, B. Harding, and A. Vince, *The entropy of an overlapping dynamical system*, arXiv:1110.6692.

[2] M. Barnsley and A. Vince, *A characaterization of the critical itineraries of an overlapping function*, arXiv:1205.5902.

[3] M. Barnsley and A. Vince, *Fractal Homeomorphism for Bi-affine Iterated Function Systems*, arXiv:1110.4680.

[4] G. Bergman, *A Number System with an Irrational Base*, Mathematics Magazine **31** (1957) 98-110.

[5] D. E. Knuth, *An imaginary number system*, Comm. ACM, **3** (1960) 245-247.

[6] D. E. Knuth, *The Art of Computer Programming, Vol. 2: Seminumerical Algorithms*, Addison-Wesley, Reading, Mass., 1969.

[7] W. Parry, *On the β-expansions of real numbers*, Acta Math. Acad. Sci. Hungar. **11** (1960) 401-416.

[8] A. Rényi, *Representations for real numbers and their ergodic properties*, Acta Math. Acad. Sci. Hungar, **8** (1957) 477-493.

[9] T. Samuel, N. Snigireva, A. Vince, *An algorithm for computing the entropy of an overlapping dynamical system*, preprint.

[10] A. Vince, *Digit tiling of Euclidean space*, in Directions in Mathematical Quasicrystals, M. Baake, R. Moody, eds. (2000) 329-370.